Seat No.:

Enrolment No.

[07]

GUJARAT TECHNOLOGICAL UNIVERSITY BE - SEMESTER-V • EXAMINATION – SUMMER 2013

Subject Code: 150503Date: 21-05-2013Subject Name: Chemical Engineering Thermodynamics – IITime: 10.30 am - 01.00 pmTotal Marks: 70Instructions:

1. Attempt all questions.

- 2. Make suitable assumptions wherever necessary.
- 3. Figures to the right indicate full marks.

Q ó 1

(a) State Lewis - Randall rule and Henryøs law. Derive the mathematical expression for fugacity and fugacity coefficient. [07]

(b) Explain in brief: ideal solutions and non-ideal solutions.

Q ó 2

(a) Discuss the various form of Gibbs-Duhem equations in detail. [07]

(b) Explain effect of temperature, pressure and total stoichiometric coefficient on equilibrium constant. Also discuss relation of equilibrium constants to composition. [07]

OR

(b) Define azeotropes and explain the minimum boiling and maximum boiling azeotropes with suitable examples. [07]

Q ó 3

(a) Explain the equilibrium conversion in heterogeneous system and multireaction system with at least one example. [07]

(b) Mention different activity coefficient model equations. Explain any one of them in detail. [07]

OR

Q ó 3

(a) Explain the reaction coordinate and its physical significance. Explain the different factors affecting equilibrium conversions. [07]

(b) Derive the expression of vapor composition at equilibrium using flash vaporization [07]

Q ó 4

(a) Calculate standard Gibbs free energy change and equilibrium constant at 700 K for the reaction:

$$N_{2(g)} + 3 H_{2(g)} \rightarrow 2 NH_{3(g)}$$

For given reaction: $\Delta H^0 = -92000 \text{ J} / \text{mol}$ and $\Delta G^0 = -33000 \text{ J} / \text{mol}$. The standard heat of formation and free energy of NH₃ at 298 K is - 46000 J / mol and - 16500 J / mol respectively. Specific heat for nitrogen, hydrogen and ammonia is given below and should be in J/mol K.

 $C_{P, N2} = 27.27 + 4.93 \text{ x } 10^{-3} \text{ T}, C_{P, H2} = 27.01 + 3.51 \text{ x } 10^{-3} \text{ T} \text{ and } C_{P, NH3} = 29.75 + 25.11 \text{ x } 10^{-3} \text{ T}$ [07]

(b) The excess Gibbs energy of a particular ternary liquid mixture is represented by the empirical expression with parameters A_{12} , A_{13} , and A_{23} functions of T and P only:

$$\frac{G^E}{RT} = A_{12} x_1 x_2 + A_{13} x_1 x_3 + A_{23} x_2 x_3$$

(i) Determine the implied expressions for \ln_1 , \ln_2 , \ln_3 .

(ii) Determines expressions (or values) for \ln_{-1} for the limiting cases: $X_1 = 0$, $x_1 = 1$, $x_2 = 0$, and $x_3 = 0$. [07]

OR

Q ó 4 For the system methanol and methyl acetate, the following equation provide reasonable correlation for activity coefficient. $\ln \gamma_1 = A X_2^2$ $\ln \gamma_2 = A X_1^2$

$$\ln P_1^{\text{sat}} = 16.591 - \frac{3643.31}{T - 33.424} \qquad \qquad \ln P_2^{\text{sat}} = 14.253 - \frac{2665.54}{T - 53.424}$$

(i) Calculate T and $\{y_i\}$ for P= 101.33 kPa, $X_1 = 0.85$

(ii) Calculate T and $\{X_i\}$ for P= 101.33 kPa, $y_1 = 0.40$

Q ó 5

(a) Discuss the application of equilibrium criteria to chemical reaction and develop expressions for the mole fractions of reacting species as functions of the reaction coordinate for:

A system initially contains 2-mol NH₃ and 5-mol O₂ undergoing the reaction:

$$4NH_{3(g)} + 5O_{2(g)} \rightarrow 4NO_{(g)} + 6H_2O_{(g)}$$

A system initially contains 3-mol H₂S and 5-mol O₂ undergoing the reaction:

$$2H_2S_{(g)} + 3O_2_{(g)} \rightarrow 2H_2O_{(g)} + 2SO_2_{(g)}$$
 [07]

[14]

(b) In a laboratory 30 mol % of methanol $\acute{0}$ water solution is to be prepared. How many m^3 of pure methanol and pure water are to be mixed to prepare 2 m^3 of desired solution?

Partial molar properties of methanol and water are given below.

Methanol: $\overline{V_1} = 38.632 \text{ cm}^3 / \text{ mol}$, Water: $\overline{V_2} = 17.765 \text{ cm}^3 / \text{ mol}$.

For the pure species at 25 0 C, methanol: V₁ = 40.227 cm³/ mol and water: V₂ = 18.068 cm³/ mol. [07]

OR

Q ó 5

(a) Using the fundamental properties relation for single phase reaction, show that [07]

$$\Delta G^0 = \circ RT \ln K$$

(b) The ammonia synthesis reaction written as: $0.5 N_{2(g)} + 1.5 H_{2(g)} \rightarrow NH_{3(g)}$ with 0.5 mol nitrogen and 1.5 mol hydrogen as the initial amounts of reactants and with the assumption that the equilibrium mixture is an ideal gas, show that: [07]

$$\varepsilon_{e} = 1 \circ (1 + 1.299 \text{KP})^{\circ 0.5}$$
