## GUJARAT TECHNOLOGICAL UNIVERSITY

BE SEMESTER 1st / 2nd (NEW) EXAMINATION WINTER 2016

Subject Code: 2110015 Date:20/01/2017

**Subject Name: Vector Calculus and Linear Algebra** 

Time:10:30 AM TO 1:30 PM **Total Marks: 70** 

**Instructions:** 

1. Question No. 1 is compulsory. Attempt any four out of remaining Six questions.

- 2. Make suitable assumptions wherever necessary.
- 3. Figures to the right indicate full marks.

## **Q.1 Objective Question (MCQ)**

Mark

(a) Choose the appropriate answer for the following MCQs.

**07** 

- 1. Rank of 3 × 3 invertible matrix is
  - a) 1
- b) 2
- c) 3
- d) 4
- Let A be a Skew-Hermitian matrix then  $A = \underline{\hspace{1cm}}$ 

  - a)  $A^T$  b)  $(\overline{A})^T$  c)  $-A^T$
- d)  $-(\overline{A})^T$
- The set  $S = \{1, x, x^2, x^3\}$  span which of the following?

- a)  $P_3$  b) R c)  $R^3$
- Let u = (1, -2) be a vector in  $\mathbb{R}^2$  with the Euclidean inner product, then ||u|| is
  - a) 1
- b) 5
- c)  $\sqrt{5}$  d)  $\sqrt{3}$
- If A is a matrix with 6 columns and rank(A) = 2, then Nullity(A) is
  - a) 2
- b) 4
- c) 0
- d) None of these
- If  $\lambda_1 = 2$ ,  $\lambda_2 = 6$  are the eigen values of the matrix A, then the eigen values of  $A^{T}$  are
  - a) 2 & 6

- b)  $\frac{1}{2} \& \frac{1}{6}$  c) 4 & 36 e) None of these
- The product of the eigen values of matrix  $A = \begin{bmatrix} 4 & 2 & -2 \\ -5 & 3 & 2 \\ -2 & 4 & 1 \end{bmatrix}$  is,

<b>(b)</b>	<b>Choose the</b>	appropriate	answer for	the follo	owing MCQs.
(~)	0110000 1110	app-op-in-			5 11 B C & 51

Let  $I:V \to V$  be an identity operator, then  $\ker(I)$  is,

a) *v* 

**b)** {0}

c) 0

d) None of these

2. The mapping  $T: \mathbb{R}^3 \to \mathbb{R}^3$  defined by T(x, y, z) = (x, y, 0) is called as

a) Projection

b) Reflection

c) Rotation

d) Magnification

If  $f_1 = x$  and  $f_2 = \sin x$ , then Wronskian  $W\left(\frac{\pi}{2}\right) =$ 

a)  $\frac{\pi}{-}$ 

b) 1

c) 0

d) -1

If  $\overrightarrow{r} = xi + yj + zk$ , then divergence of  $\overrightarrow{r}$  is

d) -3

5. The value of  $curl(grad \phi)$ , where  $\phi = 2x^2 - 3y^2 + 4z^2$  is

a) 4xi - 6yj + 8zk b) 4x - 6y + 8z c) 6

**d)** 0

6. The weighted Euclidean inner product  $\langle u, v \rangle = 3u_1v_1 + 2u_2v_2$  is the inner product on  $R^2$  generated by

a)  $\begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$  b)  $\begin{bmatrix} \sqrt{3} & 0 \\ 0 & \sqrt{2} \end{bmatrix}$  c)  $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$  d)  $\begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix}$ 

If v is a finite-dimensional vector space, and  $T: V \to V$  is a linear operator and  $ker(T) = \{0\}$ , then

a)  $R(T) \neq V$ 

b) T is one-to-one

c) Nullity  $(T) \neq 0$ 

d)None of these

Convert the following matrix in to reduced row echelon form and hence find **Q.2** 03 the rank of a matrix.

 $A = \begin{vmatrix} 1 & 4 & 2 \end{vmatrix}$ 

Solve the following system of equations by Gauss Elimination method 04  $x_1 - 2x_2 + 3x_3 = -2$ 

 $-x_1 + x_2 - 2x_3 = 3$ 

 $2x_1 - x_2 + 3x_3 = -7$ 

Method

(ii) For what choices of parameter the following system is consistent.

$$x_1 + x_2 + 2x_3 + x_4 = 1$$

$$x_1 + 2x_3 = 0$$

$$2x_1 + 2x_2 + 3x_3 = \lambda$$

- $x_2 + x_3 + 3x_4 = 2\lambda$
- **Q.3** (a) Consider the vectors u = (1, 2, -1) and v = (6, 4, 2) in  $\mathbb{R}^3$ . Show that w = (9, 2, 7) is a linear combination of u and v.
  - (b) Determine a basis for and the dimension of the solution space of the homogeneous system

$$2x_{1} + 2x_{2} - x_{3} + x_{5} = 0$$

$$-x_{1} - x_{2} + 2x_{3} - 3x_{4} + x_{5} = 0$$

$$x_{1} + x_{2} - 2x_{3} - x_{5} = 0$$

$$x_{3} + x_{4} + x_{5} = 0$$

- (c) Show that the set of all pairs of real numbers of the form (1, x) with the operations (1, y) + (1, y') = (1, y + y') and k(1, y) = (1, ky) is a vector space.
- Q.4 (a) Sketch the unit circle in an xy coordinate system in  $R^2$  using

  1. The Euclidean inner product  $\langle u, v \rangle = u_1v_1 + u_2v_2$ .
  - 2. The weighted Euclidean inner product  $\langle u, v \rangle = \frac{1}{9}u_1v_1 + \frac{1}{4}u_2v_2$ .
  - **(b)** Attempt the following.
    - 1. Let  $u=(u_1,u_2)$  and  $v=(v_1,v_2)$  be vectors in  $\mathbb{R}^2$ . Verify that the weighted Euclidean inner product  $\langle u,v\rangle=3u_1v_1+2u_2v_2$  satisfies the four inner product axioms.

04

07

- 2. Let  $R^4$  have the Euclidean inner product. Find the cosine of the angle  $\theta$  between the vectors u = (4,3,1,-2) and v = (-2,1,2,3).
- Consider the vector space  $R^3$  with the Euclidean inner product. Apply the Gram-Schmidt process to transform the basis vectors  $u_1 = (1,1,1), u_2 = (0,1,1)$  &  $u_3 = (0,0,1)$  into an orthogonal basis  $\{v_1, v_2, v_3\}$ ; then normalize the orthogonal basis vectors to obtain an orthonormal basis  $\{q_1, q_2, q_3\}$ .
- Q.5 (a) Let  $T: P_1 \to P_2$  be the linear transformation defined by T(p(x)) = x(p(x)). Find the matrix for T with respect to the standard bases  $B = \{u_1, u_2\}$  and  $B' = \{v_1, v_2, v_3\}$ , where  $u_1 = 1$ ,  $u_2 = x$ ;  $v_1 = 1$ ,  $v_2 = x$ ,  $v_3 = x^2$ .
  - (b)  $\begin{bmatrix} 4+2i & 7 & 3-i \\ 0 & 3i & -2 \\ 5+3i & -7+i & 9+6i \end{bmatrix}$  as the sum of a Hermitian and a skew-

Hermitian matrix.

(c) State the Dimension theorem for Linear Transformation and find the rank and nullity

of  $T_A$ , where  $T_A: R^6 \to R^4$  be multiplication by

$$A = \begin{bmatrix} -1 & 2 & 0 & 4 & 5 & -3 \\ 3 & -7 & 2 & 0 & 1 & 4 \\ 2 & -5 & 2 & 4 & 6 & 1 \\ 4 & -9 & 2 & -4 & -4 & 7 \end{bmatrix}$$

- - For the matrix  $A = \begin{bmatrix} -4 & -6 \\ 3 & 5 \end{bmatrix}$ , find the nonsingular matrix P and the diagonal matrix P such that  $D = P^{-1}AP$ .
  - (c) Determine  $A^{-1}$  by using Cayley-Hamilton theorem for the matrix  $A = \begin{bmatrix} 1 & 3 & 2 \\ 0 & -1 & 4 \\ -2 & 1 & 5 \end{bmatrix}$ . Hence find the matrix represented by
- Q.7 (a) Show that  $F = (y^2 z^2 + 3yz 2x)i + (3xz + 2xy)j + (3xy 2xz + 2z)k$  is both solenoidal and irrotational.
  - Find the work done when a force  $F = (x^2 y^2 + 2x)i (2xy + y)j$  moves a particle in the xy-plane from (0,0) to (1,1) along the parabola  $y^2 = x$ . Is the work done different when the path is the straight line y = x?
  - (c) State Green's theorem and use it to evaluate the integral  $\iint_C y^2 dx + x^2 dy$ , where C is the triangle bounded by x = 0, x + y = 1, y = 0.