Seat No.:	Enrolment No

Subject Name: Computer Oriented Statistical Methods

Subject Code: 151601

Instructions:

Time: 10.30 am - 01.00 pm

GUJARAT TECHNOLOGICAL UNIVERSITY

BE - SEMESTER-V • EXAMINATION - WINTER 2013

Date: 27-11-2013

Total Marks: 70

	1. 2. 3.	Mak	empt all questions. Ke suitable assumptions wherever necessary. Bures to the right indicate full marks.						
Q.1	(a)	(i)	Define normal form of a floating point number. Also, prove that for three floating point numbers a, b and c, $a \times \frac{b}{c}$ may not be equal to $\frac{a}{c} \times b$.	04					
		(ii)	For $a = 3.141592$ and an approximate value of a is 3.14, evaluate absolute error, relative error and percentage error.						
	(b)	Evaluate $\int_{-1}^{1} \frac{1}{1+x^2} dx$ by one point and two points Gaussian quadrature formula. Compare your answer with the exact value of the integral.	04						
		Explain ill-conditioned system of the linear equations.	03						
Q.2	(a)		Perform three iterations of the Newton-Raphson method to solve the system of equations $x^2 + xy + y^2 = 7$, $x^3 + y^3 = 9$.						
	(b)	(i)	Take the initial approximations as $x_0 = 1.5$, $y_0 = 0.5$. Define rate of convergence and prove that the Newton-Raphson method has second order convergence.						
		(ii)	Perform two operations of the bisection method to find an approximate root of $f(x) = 3x^3 + 10x^2 + 10x + 7 = 0$.	03					
	<i>a</i> >		OR	04					
	(b)	(i) State Budan's theorem and apply it to find the number of roots of the equation $f(x) = x^4 - 4x^3 + 3x^2 - 10x + 8 = 0$ in the interval $[-1,0]$.							
		(ii)	State Descarte's rule of signs and apply it to $P_5(x) = 8x^5 + 12x^4 - 10x^3 + 17x + 5 = 0.$	03					
Q.3	(a)		Obtain the cubic spline approximation for the function given by the data	07					
			$\begin{array}{c ccccccccccccccccccccccccccccccccccc$						
	(b)		With $M(0) = 0$, $M(3) = 0$. Hence find an estimate of $f(2.5)$. Find the linear polynomial approximation to the function $x^3 + 2$ on [1,2] OR	07					
Q.3	(a)		Consider the following table:	07					
			$\begin{array}{ c c c c c c c c c c c c c c c c c c c$						
			Find the value of x where $f(x) = 0.399$ using inverse interpolation.						
	(b)	,							
	(2)		x 1891 1901 1911 1921 1931						
			f(x) 46 66 81 93 101						
Q.4	(a)		State Trapezoidal rule. Evaluate $\int_{0.2}^{1.4} (\sin x - \log x + e^x) dx$ with $h = 0.1$ by	07					
			Simpson's $\frac{1}{3}$ and $\frac{3}{8}$ rules.						
	(b)	(i)							

1

(ii) Using Taylor series method, find y(1.1) correct to four decimal places, given 03 that $\frac{dy}{dx} = xy^{\frac{1}{2}}$, y(1) = 1

OR

Q.4 (a) Solve the following system by Gauss-Jacobi method:

27x + 6y - Z = 85, 6x + 15y + 2z = 72, x + y + 54z = 110

- (b) Apply fourth order Runge-Kutta method to find an approximate value of y for x = 0.2 with h = 0.1 if $\frac{dy}{dx} = x + y^2$, given that y = 1 when x = 0.
- Q.5 (a) (b) Obtain the median for the following distribution:

 x
 1
 2
 3
 4
 5
 6
 7
 8
 9

 f
 8
 10
 11
 16
 20
 25
 15
 9
 6

- (ii) Three unbiased coins are tossed, then find the expectation of number of tails. 03

OR

Q.5 (a) Obtain the median for the following distribution:

Mid value: Frequency: Cummulative f:

(b) The probability distribution of a random variable x is given below

Tare products	ing distinction of a range of the state of t					
x	-2	-1	0	1	2	
p(x)	1	1	p	1	1	
	12	3		$\frac{\overline{4}}{4}$	6	

Find (i) E(x), (ii) E(2x + 3), (iii) $E(x^2 + 2)$

7

7

7