GUJARAT TECHNOLOGICAL UNIVERSITY MCA - SEMESTER-II • EXAMINATION – SUMMER • 2014

Subject Code: 620005Date: 24-06-2014Subject Name: Computer Oriented Numerical MethodsTime: 10:30 am - 01:00 pmInstructions:

- 1. Attempt all questions.
- 2. Make suitable assumptions wherever necessary.
- 3. Figures to the right indicate full marks.
- Q.1 (a) What are the sources of errors? Explain any two in detail. 07
 - (b) Discuss characteristics of numerical computation with example(s).
- **Q.2** (a) Use Newton-Raphson method to find a real root of the equation $x^3 2x 5 = 0$ correct 07 to three decimal places.
 - (b) Use Bisection method in maximum six stages to find a real root of the equation $f(x) = 07 \cos x x e^x = 0$.

OR

- (b) Use Bisection method in four stages to find a real root of the equation $f(x) = x \log_{10x} 07$ 102 = 0.
- Q.3 (a) Evaluate the values of f(2) and f(6.3) using Lagrangian interpolation formula for the 07 table of values given below.

Х	1.2	2.5	4	5.1	6	6.5
f(x)	6.84	14.25	27	39.21	51	58.25

(b) Compute the value of f(7.5), by using suitable interpolation on the following table of 07 data.

Х	3	4	5	6	7	8	
f(x)	28	65	126	217	344	513	
OR							

Q.3 (a) Use the method of least square approximation to fit a straight line to the following 07 observed data.

Xi		60	61	62	63	64
Yi	i	40	40	48	52	55

(b) The following table of x and y is given

	Х	1	2	3	4		
	у	1.5	2.2	3.1	4.3		
1.							

Use cubic spline interpolation to compute y(1.2) and y'(1).

- Q.4 (a) Show that the matrix $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, satisfies the Matrix equation $A^2 - 4A - 5I = 0$. Hence find A^{-1} .
 - (b) Solve the system of equations x1 + x2 + x3 = 6 2x1 + x2 + 3x3 = 133x1 + 3x2 + 4x3 = 20

by the Gauss elimination method.

07

07

07

07

Q.4 (a) The population of a city is given in the following table. Find the rate of growth in 07 population in the year 2001 and in 1995.

Year x	1961	1971	1981	1991	2001
Population y	40.62	60.80	79.95	103.56	132.65

(b)

Evaluate $\int_{e}^{1} e^{-x^2} dx$, using

0

(1) Simpson's one-third rule with 10 sub-intervals and

(2) Trapezoidal rule.

Q.5 (a) Compute the largest Eigen value and the corresponding $\$

Eigenvector of the matrix $\begin{bmatrix} 1 & 3 & -1 \\ 3 & 2 & 4 \\ -1 & 4 & 10 \end{bmatrix}$ by power Method correct to two decimal digits.

(b) Given $xy = x - y^2$, y(2) = 1, evaluate y(2.1), y(2.2) and y(2.3) correct to four decimal **07** places using Taylor series method.

OR

- Q.5 (a) Compute values of y(0.1) and y(0.2) by 4th order Runge-Kutta method, correct to five 07 significant figures for the initial value problem. dy/dx = x + y, y(0) = 1.
 - (b) For the initial value problem, dy/dx = xy + 1, y(0) = 1, y(0.1) = 1.1053, y(0.2) = 071.22288, y(0.3) = 1.35526. Compute y(0.4) using Milne's predictor-corrector method.

07

07