GUJARAT TECHNOLOGICAL UNIVERSITY BE / ME / MBA / MCA - SEMESTER- • EXAMINATION – WINTER 2014

Subject Code: 2710710 Subject Name: LINEAR ALGEBRA Time:

Instructions:

- 1. Attempt all questions.
- 2. Make suitable assumptions wherever necessary.
- 3. Figures to the right indicate full marks.
- Q.1 (a) Are the vectors 03 $\alpha_1 = (1,1,2,4), \quad \alpha_2 = (2,-1,-5,2), \quad \alpha_3 = (1,-1,-4,0), \quad \alpha_4 = (2,1,1,6)$ linearly independent in R^4 ?
 - (b) Check whether the following set of vectors form a subspace.
 - (i) $S = \{ \alpha / \alpha = (a_1, a_2, \dots, a_n); a_1 a_2 = 0; a_i \in \mathbb{R}, \forall i \}$
 - (ii) Set of all functions of the form $k_1 + k_2 \sin x$, in $F(-\infty, \infty)$.
 - (c) Let V be a vector space over the field F. Show that the intersection of two subspaces of V is a subspace of V. What can you say for the union of two subspaces of V?
- Q.2 (a) Let V and W be two vector spaces over the field F. Let T and U be linear 07 transformations from V into W. Then prove that T + U and TU are also linear transformations.
 - (b) If T is linear transformation defined by T(1,0,0) = (0,1,0,2)

$$T(0,1,0) = (0,1,1,0)$$

$$T(0,0,1) = (0,1,-1,4)$$

Then find range of T, null space of T and dimensions of R(T) and N(T).

OR

(b) Let T be the linear operator on R^3 defined by $T(x_1, x_2, x_3) = (3x_1, x_1 - x_2, 2x_1 + x_2 + x_3).$ Is T invertible ? If so, find a rule for T^{-1} .

Q.3 (a) Let C be the field of complex numbers in algebra of polynomials and let $f = x^2 + 2$, then

- 1. If $z \in C$; where $z = \frac{1+i}{1-i}$, then show that f(z) = 1.
- 2. Find f(B), where $B = \begin{bmatrix} 1 & 0 \\ -1 & 2 \end{bmatrix}$.
- 3. If the linear operator T defined by $T(c_1, c_2, c_3) = (i\sqrt{2} c_1, c_2, i\sqrt{2} c_3)$, then find f(T).
- 4. If $g = x^4 + 3i$ then find f(g).
- (b) State and prove Cayley-Hamilton theorem.

Verify this result for $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{bmatrix}$

Date:06/01/ 2015

Total Marks: 70

04

07

07

07

07

OR

Q.3 (a) Determine the eigen values and the corresponding eigen spaces for the matrix 07 $A = \begin{bmatrix} 3 & 2 & 4 \\ 2 & 0 & 2 \\ 4 & 2 & 3 \end{bmatrix}$ (b) Let S_1 , S_2 & S_3 be the subspaces of R^3 generated by 07 $S_1 = \{(a,b,c)/a + b - 2c = 0\}, S_2 = \{(a,b,c)/a = b\}$ & $S_3 = \{(a,b,c)/a + 2c = 0, b - 4c = 0\}.$ Determine (a) a basis for $S_1 \cap S_2$ and (b) the dimension of $S_1 + S_3$. Q.4 (a) Define the following terms (i) T - invariant subspace 07

2.4 (a) Define the following terms (i)
$$T$$
 - invariant subspace
(ii) Restriction of a linear transformation

Let *T* be a linear transformation on *R*³ defined by

$$T(x, y, z) = (-3x + 3y - 2z, -7x + 6y - 3z, x - y - 2z).$$

(*i*) Show that the subspace $S = \left\{ \begin{bmatrix} 1\\2\\1 \end{bmatrix}, \begin{bmatrix} 1\\1\\-1 \end{bmatrix} \right\}$ is *T* - invariant

- (*ii*) Determine the matrix of T restricted to S with respect to the basis vectors of S.
- (b) Let R^3 have the Euclidean inner product. Use the Gram-Schmidt process to 07 transform the basis $\{u_1, u_2, u_3\}$ into an orthonormal basis where $u_1 = (0, 1, 2), \ u_2 = (-1, 0, 1), \ u_3 = (-1, 1, 3).$

OR

Q.4 (a) Let *T* be the linear operator on
$$R^3$$
 which is represented in the standard ordered 07
basis by the matrix $A = \begin{bmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{bmatrix}$. Find the minimal polynomial for *T*.
(b) Verify that the matrix $A = \begin{bmatrix} \frac{1+i}{2} & \frac{1+i}{2} \\ \frac{1-i}{2} & \frac{-1+i}{2} \end{bmatrix}$ is unitary. Also find its inverse.

Q.5 (a) Show that $\langle u, v \rangle = \frac{1}{4}u_1v_1 + \frac{1}{16}u_2v_2$ forms an inner product space on R^2 . Sketch the unit circle in an *xy*-coordinate system in R^2 .

(b) Solve the following linear system using Doolittleøs method-3x + 5y + 2z = 8 07

$$8y + 2z = -7$$
$$6x + 2y + 8z = 26$$
OR

- **Q.5** (a) Let a linear transformation on C^3 be defined by 07 $Lx = (x_1 + ix_2, x_2 + ix_3, x_3 + ix_1)$ where $x = (x_1, x_2, x_3)$ is in C^3 . Determine the adjoint of L. Show that L^* is transposed conjugate of L.
 - (b) Apply the power method to the following symmetric matrix to obtain the 07

2

dominant eigen value by taking the initial approximation as $x_0 = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^T$.

 $A = \begin{bmatrix} 0.49 & 0.02 & 0.22 \\ 0.02 & 0.28 & 0.20 \\ 0.22 & 0.20 & .040 \end{bmatrix}$
